

# Establish<sup>®</sup>

liquid

Plant Signal Technology for pea and lentil

## SEED TREATMENT COMPATIBILITY

### PEA PLANTING WINDOW

SEED TREATMENT	APPLICATION METHOD		
	SEQUENTIAL	SIMULTANEOUS	TANK MIX
Insure <sup>®</sup> Pulse	Not Recommended	Not Recommended	Not Recommended
Cruiser Maxx <sup>®</sup> Vibrance <sup>®</sup> Pulses	Not Recommended	Not Recommended	Not Recommended
Apron Maxx <sup>®</sup> RTA <sup>®</sup>	4 hours	4 hours	Not Recommended
Trilex <sup>®</sup> EverGol <sup>®</sup> + Stress Shield <sup>®</sup> (104 ml/100 kg)	Not Recommended	Not Recommended	Not Recommended
Consensus <sup>®</sup> L	24 hours	24 hours	Not Recommended
Intego <sup>™</sup> Solo	24 hours	24 hours	24 hours
Stress Shield <sup>®</sup> 600 (104 ml/100 kg)	Not Recommended	Not Recommended	Not Recommended
Cruiser <sup>®</sup> 5FS	24 hours	24 hours	24 hours
Apron XL <sup>®</sup>	Not Recommended	Not Recommended	Not Recommended

### LENTIL PLANTING WINDOW

SEED TREATMENT	APPLICATION METHOD		
	SEQUENTIAL	SIMULTANEOUS	TANK MIX
Insure <sup>®</sup> Plulse	Not Recommended	Not Recommended	Not Recommended
Cruiser Maxx <sup>®</sup> Vibrance <sup>®</sup> Pulses	Not Recommended	Not Recommended	Not Recommended
Apron Maxx <sup>®</sup> RTA <sup>®</sup>	4 hours	4 hours	Not Recommended
Trilex <sup>®</sup> EverGol <sup>®</sup> + Stress Shield <sup>®</sup> (104 ml/100 kg)	Not Recommended	Not Recommended	Not Recommended
Consensus <sup>®</sup> L	6 hours	6 hours	Not Recommended
Intego <sup>™</sup> Solo	Not Recommended	Not Recommended	Not Recommended
Stress Shield <sup>®</sup> 600 (104 ml/100 kg)	6 hours	6 hours	Not Recommended
Cruiser <sup>®</sup> 5FS	6 hours	6 hours	6 hours
Apron XL <sup>®</sup>	Not Recommended	Not Recommended	Not Recommended

### APPLICATION METHODS

**Sequential application** – seed treatment must be applied and allowed to dry before the inoculant is applied.

**Simultaneous application** – seed treatment can be applied at the same time as the inoculant but separate applicators must be used.

**Tank-mix application** – inoculant can be mixed directly with the seed treatment and applied as one solution, using one applicator. When tank mixing an inoculant with another seed treatment, dissolve the inoculant in water prior to mixing with the chemical seed treatment.

The inoculant can be tank mixed with the seed treatment for a maximum of six hours before applying to the seed. Please refer to the extended label for all other use instructions.