TECH SHEET



Imazamox and Imazethapyr herbicide Group 2

692 g (8 x 86.5 g)

Clearfield® Canola	2 to 6 leaf
Clearfield lentils	2 to 6 leaf
Clearfield sunflowers	2 to 8 leaf
Faba beans	1 to 6 leaf
Fenugreek	1 to 4 true leaf
Field Peas	1 to 6 true leaf
Seedling clover (seed production)	1 to 4 leaf
Soybeans	1 to 3 true leaf

Weeds Controlled Broadleafs

Chickweed
Cleavers
Cow Cockle
Flixweed
Hemp-nettle
Lamb's quarters
Redroot pigweed
Round-leaved mallow
Russian thistle
Shepherd's purse
Stinkweed
Volunteer canola
Wild Buckwheat

Wild mustard

Staging

Cotyledon to 4 leaf (except where indicated) (1 to 4 whorls)

Grasses

Barnyard grass 1 to 4 true leaf Green foxtail Persian darnel Volunteer barley Volunteer spring wheat Volunteer tame oats Wild oats

Application rates

Duet herbicide 17 g/ac (43 g/ha)
Adjuvant Merge® surfactant 0.5% v/v

or MSO Concentrate with Leci-Tech®

1.0% v/v

Water Volume

Ground application only 20-40 L/ac (5-10 gal/ac)

Mixing order

- 1. Use a 50 mesh (or coarser) filter screen and fill tank 3/4 full with correct amount of water. Start and continue agitation throughout mixing and spraying.
- Add the required amount of Duet herbicide watersoluble bags directly into the sprayer through the tank opening and allow to dissolve completely.
- Agitate for at least 10 minutes to dissolve the herbicide. If using a tank-mix, add the correct amount of second herbicide and continue agitating.
- 4. Continue Agitation while adding the required amount of MSO® or Merge® adjuvant. If excess foaming occurs, a silicone anti-foaming agent may be added.
- 5. Complete filling the tank to the desired level with water. If agitation is stopped for more than 5 minutes, re-suspend spray solution by full agitation prior to commencing spraying again.
- 6. Between loads of tank-mix, check in-line and nozzle screens and rinse clean if needed.





Application tips

Rainfastness - 3 hours.

Avoid application immediately before or after frost or during unseasonably cold weather.

Treat when weeds are actively growing.

Use higher water volume on dense weeds and thicker canopies.

Pre-harvest interval

60 days after application for Clearfield canola, field peas, fenugreek, faba beans and Clearfield lentils. 85 days after application for soybeans.

Follow crops

3 months after application: Winter wheat as a rotational crop

1 year after application:

Canary seed
Canola (all types incl. Clearfield canola)
Chickpeas

Durum wheat

2 years after application: Mustard (condiment type only)

Tank mixes

Please refer to herbicide label for approved tank mixes.